

University of Groningen

Mitochondrial DNA from *neurospora crassa*. Structure and genetic function

Terpstra, Peter

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1977

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Terpstra, P. (1977). *Mitochondrial DNA from neurospora crassa. Structure and genetic function*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

S U M M A R Y

The mitochondrial DNA (mtDNA) from the ascomycete *Neurospora crassa* consists of double stranded covalently closed circular molecules, with a homogeneous length of 19 μ m, as has been demonstrated by others.

We have investigated the mtDNA of wildtype strain E 5297. Mitochondria of this strain contain a population of circular DNA molecules heterogeneous in length and smaller than the 19 μ m circular DNA, which is also present in strain 5297. The buoyant density in CsCl of a covalently closed circular DNA fraction, isolated from mitochondria of strain 5297, is mitochondrial. This suggests that the small heterogeneous circular DNA could be derived from the 19 μ m circular DNA. This relationship, however, could not be investigated, because the isolation of the small circular DNA was not reproducible (Chapter 3).

mtDNA from *Neurospora crassa* strain E 5256 (19 μ m circles) was isolated on preparative scale as a collection of linear fragments. With equilibrium density centrifugation in CsCl, sedimentation analysis and denaturation-renaturation techniques it could be established that the distribution of GC basepairs in the mtDNA molecule is heterogeneous. A continuous stretch with a length of at least one third of the molecule has a 6% lower GC content than the 40% average GC content of the whole molecule (Chapter 4).

Cleavage of the mtDNA with restriction endonucleases yields a collection of specific fragments, which can be separated according to size by gel electrophoresis. Endonuclease Bam HI, Eco RI and Hind III cleave the DNA in 5, 12 and more than 30 fragments respectively. Analysis of partial cleavage products allowed us to determine the sequence of the 5 Bam HI and 12 Eco RI fragments. The 5 Bam HI recognition sites could be located on the Eco RI fragments by isolation of the individual Eco RI fragments and redigestion with endonuclease Bam HI. In this way two overlapping cleavage maps could be constructed. In addition we determined the position of the largest Hind III fragment on the map (Chapter 5).

The cleavage map of the mtDNA provided a framework for the localization of genes. From DNA-RNA hybridization experiments we conclude that the mtDNA contains one gene for the large (24S) and one for the small (17S) mitochondrial ribosomal RNA. The genes are found in the region of the mtDNA, relatively rich in GC, on two neighbouring fragments. Their mutual distance is about 900 basepairs. This confirms the existence of a ribosomal RNA precursor,

which has been found previously
cleavage map provided a starting
within the largest Eco RI fragme

Hybridization of mitochondri
fragments leads to the conclusio
found in a limited area, of maxi
the 24S gene. The other tRNA ger
preliminary experiments suggest
also a gene for a ribosomal 5S R

which has been found previously by others. The place of the 24S RNA on the cleavage map provided a starting point for ordering 8 Hind III fragments within the largest Eco RI fragment (Chapter 6).

Hybridization of mitochondrial transfer RNAs (tRNA) to mtDNA restriction fragments leads to the conclusion that about 80% of the tRNA genes can be found in a limited area, of maximally 15% genome length. This area is next to the 24S gene. The other tRNA genes are scattered over the whole genome. Some preliminary experiments suggest that in the area of the ribosomal RNA genes, also a gene for a ribosomal 5S RNA can be found (Chapter 7).

omycete *Neurospora crassa* con-
ular molecules, with a homoge-
by others.

strain E 5297. Mitochondria
DNA molecules heterogeneous
NA, which is also present in
ovalently closed circular DNA
5297, is mitochondrial. This
DNA could be derived from the
could not be investigated,
was not reproducible (Chapter

(19 μ m circles) was isolated
ragments. With equilibrium
alysis and denaturation-rena-
the distribution of GC
. A continuous stretch with
as a 6% lower GC content than
(Chapter 4).

nucleases yields a collection
ording to size by gel elec-
d III cleave the DNA in 5, 12
s of partial cleavage pro-
e 5 Bam HI and 12 Eco RI
be located on the Eco RI
fragments and redigestion
pping cleavage maps could be
ion of the largest Hind III

amework for the localization
ve conclude that the mtDNA
the small (17S) mitochon-
region of the mtDNA, relati-
their mutual distance is
of a ribosomal RNA precursor,